

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/243966215>

SARC-F: A Simple Questionnaire to Rapidly Diagnose Sarcopenia

Article in *Journal of the American Medical Directors Association* · June 2013

DOI: 10.1016/j.jamda.2013.05.018 · Source: PubMed

CITATIONS

147

READS

81,343

2 authors:



Theodore K Malmstrom
Saint Louis University

112 PUBLICATIONS 4,350 CITATIONS

[SEE PROFILE](#)



John E Morley
Saint Louis University

1,477 PUBLICATIONS 76,014 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Multimodal treatment protocols for geriatric patients with balance problems [View project](#)



Sarcopenia in elders [View project](#)



JAMDA

journal homepage: www.jamda.com

Editorial

SARC-F: A Simple Questionnaire to Rapidly Diagnose Sarcopenia

Theodore K. Malmstrom PhD^{a,b}, John E. Morley MB, BCh^{b,*}^aDepartment of Neurology and Psychiatry, Saint Louis University School of Medicine, St. Louis, MO^bDivision of Geriatric Medicine, Saint Louis University School of Medicine, St. Louis, MO

Sarcopenia was originally defined as an age-associated loss of muscle mass.^{1,2} Recently a number of European and international groups have redefined sarcopenia as being a decline in muscle function (either walking speed or grip strength) associated with loss of muscle mass.^{3–5} This approach has been validated.^{6,7} Sarcopenia leads to disability, falls, and increased mortality.^{8–16} Loss of muscle strength and aerobic function are 2 of the hallmarks of frailty.^{17–21} Sarcopenia has been linked to an increased prevalence of osteoporosis, thus further increasing its propensity to produce hip fractures.^{22–27}

Although osteoporosis has been classically diagnosed by measuring bone mineral density, it has been recognized that a number of other factors play into the role of diagnosing the propensity to have a fracture.^{28,29} This is particularly true in older persons with diabetes mellitus who often have good bone mineral density but weak bones, and this is coupled with an increase in sarcopenia.^{30–33} This has led to the concept that the questions associated with the Fracture Risk Assessment Tool (FRAX) (www.shef.ac.uk/FRAX) may be sufficient to screen for osteoporosis. Two studies have confirmed that bone mineral density does not need to be measured in most cases to make or exclude the diagnosis of osteoporosis.^{34,35} As sarcopenia is much more definable by simple functional questions, this led to the concept that a simple questionnaire could be developed to diagnose sarcopenia and obviate the need for the measurement of muscle mass.

The SARC-F questionnaire has been developed as a possible rapid diagnostic test for sarcopenia.³⁶ There are 5 SARC-F components: Strength, Assistance with walking, Rise from a chair, Climb stairs and Falls (Table 1). The scores range from 0 to 10, with 0 to 2 points for each component. Our preliminary studies have suggested that a score equal to or greater than 4 is predictive of sarcopenia and poor outcomes.

The ability to rapidly diagnose sarcopenia is important, as there is increasing evidence that therapeutic interventions can improve outcomes. Among successful therapeutic outcomes are resistance exercise,^{37–39} vitamin protein supplementation,^{40–45} and possibly testosterone.^{46–48}

The authors declare no conflicts of interest.

* Address correspondence to John E. Morley, MB, BCh, Division of Geriatric Medicine, Saint Louis University School of Medicine, 1402 S. Grand Boulevard, M238, St. Louis, MO 63104.

E-mail address: morley@slu.edu (J.E. Morley).

Table 1
SARC-F Screen for Sarcopenia

Component	Question	Scoring
Strength	How much difficulty do you have in lifting and carrying 10 pounds?	None = 0 Some = 1 A lot or unable = 2
Assistance in walking	How much difficulty do you have walking across a room?	None = 0 Some = 1 A lot, use aids, or unable = 2
Rise from a chair	How much difficulty do you have transferring from a chair or bed?	None = 0 Some = 1 A lot or unable without help = 2
Climb stairs	How much difficulty do you have climbing a flight of 10 stairs?	None = 0 Some = 1 A lot or unable = 2
Falls	How many times have you fallen in the past year?	None = 0 1–3 falls = 1 4 or more falls = 2

References

- Morley JE, Baumgartner RN, Roubenoff R, et al. Sarcopenia. *J Lab Clin Med* 2001; 137:231–243.
- Bauer JM, Kaiser MJ, Sieber CC. Sarcopenia in nursing home residents. *J Am Med Dir Assoc* 2008;9:545–551.
- Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 2010;39:412–423.
- Morley JE, Abbatecola AM, Argiles JM. Society on sarcopenia, Cachexia and Wasting Disorders Trialist Workshop. Sarcopenia with limited mobility: An international consensus. *J Am Med Dir Assoc* 2011;12:403–409.
- Fielding RA, Vellas B, Evans WJ, et al. Sarcopenia: An undiagnosed condition in older adults. Current consensus definition: Prevalence, etiology, and consequences. International Working Group on Sarcopenia. *J Am Med Dir Assoc* 2011;12:249–256.
- Malmstrom TK, Miller DK, Herning MM, Morley JE. Low appendicular skeletal muscle mass (ASM) with limited mobility and poor health outcomes in middle-aged African Americans [published online ahead of print March 27, 2013]. *J Cachexia Sarcopenia Muscle*.
- Patel HP, Syddall HE, Jameson K, et al. Prevalence of sarcopenia in community-dwelling older people in the UK using the European Working Group on Sarcopenia in Older People (EWGSOP) definition: Findings from the Hertfordshire Cohort Study (HCS). *Age Ageing* 2013;42:378–384.
- Roman D, Mahoney K, Mohamadi A. Sarcopenia: What's in a name? *J Am Med Dir Assoc* 2013;14:80–82.
- Cesari M, Vellas B. Sarcopenia: A novel clinical condition or still a matter for research? *J Am Med Dir Assoc* 2012;13:766–767.
- Landi F, Liperoti R, Fusco D, et al. Sarcopenia and mortality among older nursing home residents. *J Am Med Dir Assoc* 2012;13:121–126.

11. Neyens JC, van Haastregt JC, Dijcks BP, et al. Effectiveness and implementation aspects of interventions for preventing falls in elderly people in long-term care facilities: A systematic review of RCTs. *J Am Med Dir Assoc* 2011;12:410–425.
12. Choi M, Hector M. Effectiveness of intervention programs in preventing falls: A systematic review of recent 10 years and meta-analysis. *J Am Med Dir Assoc* 2012;13:188.e13–188.e21.
13. Morley JE, Rolland Y, Tolson D, Vellas B. Increasing awareness of the factors producing falls: The mini falls assessment. *J Am Med Dir Assoc* 2012;13:87–90.
14. Vu MQ, Weintraub N, Rubenstein LZ. Falls in the nursing home: Are they preventable? *J Am Med Dir Assoc* 2006;7:S53–S58. 52.
15. Rapp K, Becker C, Cameron ID, et al. Epidemiology of falls in residential aged care: Analysis of more than 70,000 falls from residents of Bavarian nursing homes. *J Am Med Dir Assoc* 2012;13:187.e1–187.e6.
16. Arango-Lopera VE, Arroyo P, Gutiérrez-Robledo LM, et al. Mortality as an adverse outcome of sarcopenia. *J Nutr Health Aging* 2013;17:259–262.
17. Rougé Bugat ME, Cestac P, Oustric S, et al. Detecting frailty in primary care: A major challenge for primary care physicians. *J Am Med Dir Assoc* 2012;13:669–672.
18. Abellan van Kan G, Rolland YM, Morley JE, Vellas B. Frailty: Toward a clinical definition. *J Am Med Dir Assoc* 2008;9:71–72.
19. Morley JE, Vellas B, Abellan van Kan G, et al. Frailty consensus: A call to action. *J Am Med Dir Assoc* 2013;14:392–397.
20. Peters LL, Boter H, Buskens E, Slaets JP. Measurement properties of the Groningen Frailty Indicator in home-dwelling and institutionalized elderly people. *J Am Med Dir Assoc* 2012;13:546–551.
21. Gobbens RJ, van Assen MA, Luijckx KG, et al. Determinants of frailty. *J Am Med Dir Assoc* 2010;11:356–364.
22. Sjoblom S, Suuronen J, Rikkonen T, et al. Relationship between postmenopausal osteoporosis and the components of clinical sarcopenia. *Maturitas* 2013;75:175–180.
23. Cederholm T, Cruz-Jentoft AJ, Maggi S. Sarcopenia and fragility fractures. *Eur J Phys Rehabil Med* 2013;49:111–117.
24. Miyakoshi N, Hongo M, Mizutani Y, Shimada Y. Prevalence of sarcopenia in Japanese women with osteopenia and osteoporosis [published online ahead of print March 21, 2013]. *J Bone Miner Metab*.
25. Morley JE. Falls and fractures. *J Am Med Dir Assoc* 2007;8:276–278.
26. Kaji H. Linkage between muscle and bone: Common catabolic signals resulting in osteoporosis and sarcopenia. *Curr Opin Clin Nutr Metab Care* 2013;16:272–277.
27. Morley JE. Frailty, falls, and fractures. *J Am Med Dir Assoc* 2013;14:149–151.
28. Morley JE. Osteoporosis and fragility fractures. *J Am Med Dir Assoc* 2011;12:389–392.
29. Martinez-Reig M, Ahmad L, Duque G. The orthogeriatrics model of care: Systematic review of predictors of institutionalization and mortality in post-hip fracture patients and evidence for interventions. *J Am Med Dir Assoc* 2012;13:770–777.
30. Migdal A, Yarandi SS, Smiley D, Umpierrez GE. Update on diabetes in the elderly and in nursing home residents. *J Am Med Dir Assoc* 2011;12:627–632.e2.
31. Morley JE. Diabetes mellitus: “The times they are a–changing.” *J Am Med Dir Assoc* 2012;13:574–575.
32. Leenders M, Verdijk LB, van der Hoeven L, et al. Patients with type 2 diabetes show a greater decline in muscle mass, muscle strength, and functional capacity with aging. *J Am Med Dir Assoc* 2013;14:585–592.
33. Sinclair A, Morley JE, Rodriguez-Mañas L, et al. Diabetes mellitus in older people: Position statement on behalf of the International Association of Gerontology and Geriatrics (IAGG), the European Diabetes Working Party for Older People (EDWPOP), and the International Task Force of Experts in Diabetes. *J Am Med Dir Assoc* 2012;13:497–502.
34. Kanis JA, McCloskey E, Johansson H, et al. FRAX® with and without bone mineral density. *Calcif Tissue Int* 2012;90:1–13.
35. Leslie WD, Lix LM, Langsetmo L, et al. Construction of a FRAX® model for the assessment of fracture probability in Canada and implications for treatment. *Osteoporos Int* 2011;22:817–827.
36. Malmstrom TK, Morley JE. Sarcopenia: The target population. *J Frailty Aging* 2013;2:55–56.
37. Yamada M, Arai H, Sonoda T, Aoyama T. Community-based exercise program is cost-effective by preventing care and disability in Japanese frail older adults. *J Am Med Dir Assoc* 2012;13:507–511.
38. Singh NA, Quine S, Clemson LM, et al. Effects of high-intensity progressive resistance training and targeted multidisciplinary treatment of frailty on mortality and nursing home admissions after hip fracture: A randomized controlled trial. *J Am Med Dir Assoc* 2012;13:24–30.
39. Valenzuela T. Efficacy of progressive resistance training interventions in older adults in nursing homes: A systematic review. *J Am Med Dir Assoc* 2012;13:418–428.
40. Demontiero O, Herrmann M, Duque G. Supplementation with vitamin D and calcium in long-term care residents. *J Am Med Dir Assoc* 2011;12:190–194.
41. McKinney JD, Bailey BA, Garrett LH, et al. Relationship between vitamin D status and ICU outcomes in veterans. *J Am Med Dir Assoc* 2011;12:208–211.
42. Mithal A, Bonjour JP, Boonen S, et al, IOF CSA Nutrition Working Group. Impact of nutrition on muscle mass, strength, and performance in older adults. *Osteoporos Int* 2013;24:1555–1566.
43. Morley JE, Argiles JM, Evans WJ, et al. Society for Sarcopenia, Cachexia, and Wasting Disease. Nutritional recommendations for the management of sarcopenia. *J Am Med Dir Assoc* 2010;11:391–396.
44. Tieland M, van de Rest O, Dirks ML, et al. Protein supplementation improves physical performance in frail elderly people: A randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 2012;13:720–726.
45. Tieland M, Kirks ML, van der Zwaluw N, et al. Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: A randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 2012;13:713–719.
46. Morley JE. Anabolic steroids and frailty. *J Am Med Dir Assoc* 2010;11:533–536.
47. Wittert GA, Chapman IM, Haren MT, et al. Oral testosterone supplementation increases muscle and decreases fat mass in healthy elderly males with low-normal gonadal status. *J Gerontol A Biol Sci Med Sci* 2003;58:618–625.
48. Morley JE. Androgens and aging. *Maturitas* 2001;38:61–71.